



Remote Sensing and Landscape Metrics for Global Urban Ecological Monitoring

Maik Netzband¹, William L. Stefanov² ¹Center for Environmental Studies, Arizona State University, Box 873211, Tempe, AZ 85287-3211, ²Geological Sciences, Arizona State University, Tempe, AZ 85287-6305



Central Arizona Phoenix Region
ASTER - Visible and Near Infrared



Mexico City Metropolitan Area
ASTER – Visible and Near Infrared




Fig. 1




Fig. 2




Fig. 3

The 'Patch density' (fig. 1) equals the number of patches of the corresponding patch type (NP) divided by total landscape area, multiplied by 10,000 and 100 (to convert to 100 hectares). It is an index for the fragmentation of a landscape and facilitates the comparison of landscapes with different sizes. This metric shows higher values for all classes in the Phoenix area except for the 'High Vegetation'.

'Mean patch size' (fig. 2) indicates the medium patch size in hectares and is a function of the total area of the landscape and the patch number. Smaller values point to a higher fragmentation of the landscape. The 'Largest Patch Index' (fig. 3) is the average 'perimeter-to-area-ratio' for a certain class. A low value indicates a very compact form with small edge lengths, whereas higher values express more complex forms. Both indices show a similar distribution for the compared urban areas: there is a peak in the Mexico City area for the vegetation classes ('moderate' and 'high') as well as for the 'Low Density Urban' class. While peaks for this class and the 'High Vegetation' class are also present for Phoenix, a more significant peak is present for the 'Bare Soil/Low Vegetation' class.




Fig. 4




Fig. 5





Fig. 6


'Edge density' (fig. 4) equals the sum of the lengths of all edge segments involving the corresponding patch type, divided by the total landscape area, converted to hectares. Here the urban classes show higher values in Phoenix compared to Mexico City, whereas the vegetation classes are distributed vice versa.

The value of 'Patch size coefficient of variation' (fig. 5) becomes 0, if the investigated landscape consists of one (or equally large) patches. A high value displays a strong variation in the landscape sample. The variation is higher again within the urban classes of the Phoenix area, although the highest values of this index can be found in the vegetation classes of the Mexico City area.

The 'Area-weighted mean shape index' (fig. 6) calculates the complexity of the patches in the landscape and these are weighted according to their size. Larger patches enter more strongly in the calculation than smaller ones do. The most significant differences between the two compared urban agglomerations can be observed in the 'Moderate Density Urban' class (higher in Phoenix), as well as in the Moderate and High Vegetation classes which show higher values in Mexico City. This can be identified as general pattern for most of the analysed metrics. The 'Bare Soil/Low Vegetation' class is always higher in the Phoenix area.



Central Arizona Phoenix Region
ASTER – Land Cover Classification



Mexico City Metropolitan Area
ASTER – Land Cover Classification

For more information visit:

<http://elwood.la.asu.edu/grs/UEM/>

References

- Frohn, R.C. (1998). Remote sensing for landscape ecology. New metric indicators for monitoring, modeling, and assessment of ecosystems, 99 p., Boca Raton.
- Netzband, M., Kirstein, W. (2001). Landscape metrics as a tool for the comparison of different urban regions, *Regenburger Geographische Schriften* 35, pp. 113-132 (CD-ROM).
- Stefanov, W.L. (2002). Assessment of landscape fragmentation associated with urban centers using ASTER data, *American Geophysical Union EOS Transactions* 83B, 0739.
- Stefanov, W.L., Christensen, P.R., and Ramsey, M.S. (2016a). Remote sensing of urban ecology at regional and global scales: Results from the Central Arizona Phoenix Urban Environmental Monitoring program, *Regenburger Geographische Schriften* 55, pp. 113-124 (on supplemental CD-ROM).
- Stefanov, W.L., Ramsey, M.S., and Christensen, P.R. (2016b). Monitoring urban land cover change: An expert system approach to land cover classification of sensor centers, *Remote Sensing of Environment* 77 (2), pp. 173-185.